

Patient information

Bowel Preparation for Small Bowel Capsule Endoscopy (Plenvu)

Gastroenterology Department

Your clinician would like you to have bowel preparation for your small bowel capsule endoscopy. The instructions are given below and should be followed carefully.

What is Bowel Preparation and what are the benefits of taking it before the test?

Bowel preparation is medicine used to clear the waste material from the bowel in order that there are good views of the lining of the small bowel. Plenvu is a type of bowel preparation that is commonly used to clear the bowel.

This will allow for clear pictures to be taken of your small bowel lumen.

Are there any risks/side effects?

- Plenvu will clear out the bowel resulting in diarrhoea within approximately one to three hours of the first dose. It is advised that you stay within easy reach of a toilet once you start to take the medication.
- Some patients experience nausea, vomiting, or abdominal pain. This can be associated with dizziness. If any of these symptoms occur, please lie down until they pass.
- You might like to use a barrier cream to prevent a sore bottom. A product such as petroleum jelly can also be effective.
- Some stomach cramping is normal.
- If you vomit up the bowel preparation at any time or have any other concerns regarding side effects, please contact us directly – see end of leaflet for contact details

Other medication

- **Female patients: If you are taking an oral contraceptive pill, diarrhoea can make it less effective. Please use additional forms of contraception such as condoms.**
- Any other oral medication can be continued as normal unless otherwise instructed.
- **Do not take any medication two hours either side of drinking Plenvu.**
- Please bring a list of your medications that we may review this on the day.

Laxative Treatment

Three sachets of Plenvu are included in this pack. They work together to empty your bowel.

Three days before your test you must follow a low residue diet. Instructions explaining what you can eat and what you must avoid are listed further down on this leaflet.

The day before your test, you should take two doses (a total of three sachet's) of Plenvu.

How do I take the Plenvu sachets?

Dose one – at 5pm the day before your test

- Pour the contents of the single sachet for dose one into a jug.
- Make the solution up to 500mL with water.
- Stir the solution well.
- Drink the solution over a period of 60 minutes.
- Alternate sips of the solution with sips of clear fluids as outlined below.
- Aim to drink an additional 500mL of clear fluids in between sips of the Plenvu solution

Dose two (Sachet A and B) – at 9pm the day before your test

- Pour the contents of the two sachets (sachet A and B) together in a jug.
- Make the solution up to 500mL with water.
- Stir the solution well.
- Drink the solution over a period of 60 minutes.
- Alternate sips of the solution with sips of clear fluids as outlined below.
- Aim to drink an additional 500mL clear fluids in between sips of the Plenvu solution.

Diet

You must start the special diet from three days before your test. This is a low-residue diet which limits the amount of fibre and other foods that increase bowel activity.

An example of a low residue diet is shown below.

You may eat freely from the foods contained in this list, unless you are allergic or have been advised to avoid certain foods.

Please avoid taking any food or drink that is not contained within this list of foods/drink you can consume.

No solid food should be taken after 9 AM on the day before your test but you should continue to drink clear fluids.

An example of clear fluid drinks is shown further in this leaflet.

Foods you can eat

- White meat, skinless chicken, grilled/boiled/poached or steamed white fish.
- Potato without skin boiled/mashed/chips.
- Eggs, cheese, tofu.
- White - bread, croissants, pasta, rice, cous cous, pastry (white flour).
- Cornflakes, Rice Krispies, Frosted Flakes.
- Smooth yoghurts.
- Crisps.
- Butter/margarine.
- Seedless/shredless marmalade or jam.
- Marmite, honey, rich tea biscuits, chocolate, yoghurt.
- Plain ice cream, custard, clear jelly (no red or blackcurrant jelly), boiled sweets.
- Sponge cakes without fruit or nuts, rice pudding/semolina.

Foods you should avoid

- Wholemeal or granary bread, wholemeal flour, wholemeal pasta, brown rice, pearl barley, bran, quinoa.
- All whole-wheat cereals e.g Branflakes, Weetabix, Shreddies etc.
- Porridge and Muesli (all containing dried fruit and nuts).
- Yogurts or cheese containing fruit or nut pieces.
- Red meat or fish that is tough or gristly, skin and bones of fish and chicken, pies, egg dishes containing vegetables.
- Vegetables, salad, baked beans, split peas, lentils, peas, sweet corn, celery, All seeds, tough skins, potato skins.
- All fruit including dried fruit, citrus fruit and berries, prunes, smoothies, fruit juices with bits.
- Avoid all nuts including coconut and almond.
- Puddings, cakes and biscuits made with wholemeal flour, dried fruits or nuts.
- Chocolate or toffee with dried fruits or nuts, Popcorn, Marzipan, Digestive biscuits.
- Marmalade with peel or jam with seeds.
- Lentil and vegetable soups, Pickles and chutneys, Horseradish, Relish.

The day before and the day of the procedure, MILK in any form should be avoided.

What can I drink?

Clear Fluids:

- Water, mineral/fizzy, fizzy drinks (including Lucozade), fruit squash (not blackcurrant), chicken stock cubes, clear soup, tea/coffee (to be taken black the day before and day of the procedure), herbal/fruit teas.
- Sweetener may be added to food/drinks.
- Oxo, Bovril, clear soup (watery without bits in it).

You must drink at least one glass of fluid from the list every hour (throughout the daytime) during the diet preparation period.


Alcohol:

We advise to minimise alcohol consumption, in line with national guidance. If alcohol consumption is desired the following can be drank.

- Lager / Beer.
- Spirits with light coloured mixer i.e., Lemonade / Tonic.
- White wine.

Alcoholic drinks to be avoided:

- Red / Rose Wine.
- Dark Mixers including Cola.
- Guinness.



Example of low residue diet: Breakfast

- One boiled or poached egg.
- One to two slices of white bread, butter or margarine or honey.
- Tea or coffee.

Midday

- Steamed or grilled white fish or chicken.
- One medium boiled potato (no skin) or two tablespoons of boiled white rice.
- Small plain jelly.

Evening

- As midday, or other foods from the low residue foods list.
- Small plain jelly.

It is important to drink plenty of fluids in-between mealtimes

Summary of the Bowel Preparation Regime:

Please follow the regime outlined below

Day	Date	Diet	Laxative Treatment
Three days before the test		Low residue diet	Nil required
Two days before the test		Low residue diet	Nil required
One day before the test		Light breakfast only and then no further food after 9 AM with clear fluids only for the rest of the day.	Dose one of Plenvu at 5pm Dose two of Plenvu (sachet A+ B) at 9pm
Day of test		You may have sips of water if required. Please do not eat any food until 2 hours after swallowing the capsule.	Nil required

What do I do on the day of the procedure?

Remember you are not allowed to eat anything until after your test.

- You may continue to take all your other usual medications as normal. **Do not take your medication one hour either side of the planned time for swallowing the capsule.**
- Do not take tablets for diabetes on the day of your procedure but **bring your medication with you.**
- If you take insulin, please take this as instructed by your diabetes nurse.
- If you normally monitor your blood sugar, please check it every two hours.
 - **If you feel dizzy, sweaty or irritable (signs of low blood glucose) take two to three teaspoons of sugar, or six glucose tablets, or one glass of Lucozade.**
 - You should feel better almost immediately, if not repeat the above treatment. If symptoms persist, contact your GP or your specialist community diabetes team.

Attend the gastroenterology department at your appointment time.

Feedback

Your feedback is important to us and helps us influence care in the future.

Following your discharge from hospital or attendance at your outpatient appointment you will receive a text asking if you would recommend our service to others. Please take the time to text back, you will not be charged for the text and can opt out at any point. Your co-operation is greatly appreciated.

Further information

Contact the Gastroenterology Unit at the Royal Liverpool Hospital
Monday – Sunday: 8am – 6pm

Telephone: 0151 706 2819 / 2726 / 2656

Text phone number: 18001 0151 706 2819 / 2726 / 2656

Author: Gastroenterology

Review date: May 2028

All Trust approved information is available on request in alternative formats, including other languages, easy read, large print, audio, Braille, moon and electronically.

يمكن توفير جميع المعلومات المتعلقة بالمرضى الموافق عليهم من قبل الائتمان المستشفى عند الطلب بصيغ أخرى، بما في ذلك لغات أخرى وبطرق تسهل قراءتها وبالحروف الطباعية الكبيرة وبالصوت وبطريقة برايل للمكفوفين وبطريقة مون والكترونياً.

所有經信托基金批准的患者資訊均可以其它格式提供，包括其它語言、易讀閱讀軟件、大字

體、音頻、盲文、穆恩體 (Moon) 盲文和電子格式，敬請索取。

در صورت تمایل می‌توانید کلیه اطلاعات تصویب شده توسط اتحادیه در رابطه با بیماران را به اشکال مختلف در دسترس داشته باشید، از جمله به زبانهای دیگر، به زبان ساده، چاپ درشت، صوت، خط مخصوص کوران، مون و بصورت روی خطی موجود است.

زانیاری پیوندیدار بهو نه‌خوشانه‌ی له‌لایه‌ن تراسته‌وه په‌سند کراون، نه‌گمر داوا بکریت له فورماته‌کانی تردا بریتی له زمانه‌کانی تر، نیزی رید (هاسان خویندنه‌وه)، چاپی گه‌وره، شریتی ده‌نگ، هیلی موون و نه‌لیکترونیکی هیه.

所有经信托基金批准的患者信息均可以其它格式提供，包括其它语言、易读阅读软件、大字
体、音频、盲文、穆恩体（Moon）盲文和电子格式，敬请索取。

Dhammaan warbixinta bukaanleyda ee Ururka ee la oggol yahay waxaa marka la codsado lagu heli karaa nuskhado kale, sida luqado kale, akhris fudud, far waaweyn, dhegeysi, farta braille ee dadka indhaha la', Moon iyo nidaam eletaroonig ah.