

# Patient information

# Outpatient Flexible Sigmoidoscopy with Radiofrequency Ablation (RFA)

# Gastroenterology and Hepatology

Your Consultant or doctor has advised you to have a Flexible Sigmoidoscopy with Radiofrequency Ablation treatment.

# What is a Flexible Sigmoidoscopy?

This is an examination of the lower bowel in which a flexible telescope is inserted into your back passage, allowing inspection of the left bowel (colon). The scope has a lens on its end, which allows the person performing the test to view the inside of your bowel on a television screen.

# What is Radiofrequency Ablation (RFA)?

This is a type of heat treatment that is given using the telescope. It uses heat from a small electrical current to cauterise the blood vessels that form after radiotherapy in the back passage (rectum). It is these blood vessels that cause the bleeding. If necessary the procedure can be repeated after a few weeks.

#### What are benefits of Radiofrequency Ablation?

The main benefit of RFA is to stop the bleeding that you are having by cauterising the blood vessels. It may also stop you from needing blood transfusions, if you do at the moment.

# What are the risks of Flexible Sigmoidoscopy and Radiofrequency Ablation?

#### **Peritonitis**

It is possible to damage the large bowel lining making a hole, but this happens rarely. This allows bowel contents to escape causing severe inflammation in the abdomen (peritonitis). This nearly always needs an operation to repair the hole. The risk of this happening is approximately one in every 5000 examinations.

#### Ulceration and stricturing

It is possible that the treatment causes ulceration before healing and occasionally narrowings (strictures) when the bowel heals.

#### Are there any alternatives to this procedure?

At present, there are very few alternative treatments for bleeding due to radiation proctopathy. Most people having treatment for bleeding via endoscopy have tried sucralfate enema treatment, which has not worked.

Alternative endoscopic treatments of argon plasma coagulation and formalin treatment are more prone to side effects of pain, bleeding, narrowings (stricture formation) and tears (perforation). Also several treatments are needed.

With RFA the whole of the back passage (rectum) can be treated in one session and as the heat treatment is not as deep, narrowings and tears are less likely to happen.

## What will happen if I don't have this procedure done?

Doing nothing may result in your bleeding continuing.

#### What anaesthetic or sedation will I be given?

The examinations are usually done with sedation. If you do have sedation, the drugs used in sedation may affect your memory or concentration for up to 24 hours. Many patients remember nothing about the procedure or even what the doctor has said to them afterwards.

A side effect of these drugs is that they can slow your breathing – this should not normally happen but sometimes patients can be oversensitive to the drug. This is the main reason we do not give high doses of these drugs. We also will give you oxygen during the test.

If you are worried about any of these risks, please speak to your consultant or a member of their team before you are due to have this treatment.

If you have sedation, you must have a friend or relative collect you from the Gastroenterology Unit and we recommend they stay with you afterwards.

### **Getting ready for your Flexible Sigmoidoscopy**

- You will need to take either an oral bowel preparation that you will have received with your appointment or be asked to have an enema preparation to clear the lower bowel. Sometimes we instruct patients about a different preparation.
- If you are taking iron tablets, please stop taking them one week before your appointment.
- Please bring a dressing gown and slippers.
- When you come to the department, please tell the doctor or nurse about any medicines you are taking, possible allergies or bad reactions you may have had, or heart valve problems requiring antibiotics.

#### Your Flexible Sigmoidoscopy procedure

- During the test, you will be asked to lie on your left side on a couch with your knees bent.
- A rectal examination will be performed with a gloved finger, which will help to relax the muscle. After this, the telescope will be lubricated and inserted into your bottom.
- Air is then blown down the telescope and into your bowel, so that it can be seen clearly.

The RFA treatment is then given.

## After your Flexible Sigmoidoscopy

You will be able to walk back to the area you came from to get dressed. You may wish to have a friend or relative call to collect you.

# If you have had sedation you must have a friend or relative collect you from the Gastroenterology Unit and for next 24 hours you must not

- Travel alone.
- Drive any vehicle e.g. car, scooter or ride a motorcycle or bicycle.
- Operate machinery (including domestic appliances such as a kettle).
- Climb ladders.
- Make important decisions, sign any business or legal documents.
- Drink alcohol.
- Return to work within 12 hours of treatment. Your general health and any medicines you are taking may increase the time you need off work.

#### You should

- Take it easy for the rest of the day, avoid strenuous activity.
- Take your medications as usual.
- Let someone else care for anyone you usually look after, such as children or elderly or sick relatives.

### Minor post procedure symptoms

You may notice some wind pains due to the air that has been put into the bowel at the time of the examination, which is done to allow the doctor/nurse to get a clear view of your bowel. This should disperse once you are up and about walking around.

You may have some spotting of blood if you have had any polyps or large pieces of tissue samples removed.

#### **Serious post procedure symptoms**

If you experience any severe pain in your neck, chest or abdomen or if you are bleeding heavily you must return to the Accident and Emergency Department and bring this, and any other information leaflet you are given with you so as to inform the staff what you have had done.

#### Results

Results of the examination will either be given to you before you leave or at your next clinic appointment. Sometimes biopsy, photography or other information is taken which may need to be assessed further and the result may not be available for a week or two. A copy of the report will be sent to your family doctor (GP).

#### **Cancellations**

If you are unable to keep this appointment, please let us know as soon as possible on

Tel: 0151 706 2656. Text phone number: 18001 0151 706 2656

We will be able to give your appointment to another patient, and arrange another for you.

#### **Transport**

- Transport has not been arranged for this appointment so it will be necessary for you to make your own arrangements.
- Parking for patients and visitors is available at the Q-Park multi-storey car park
  opposite the main hospital entrance on Prescot Street. The entrance by car is on
  Epworth Street off Erskine Street. The car park is open 24 hours a day, seven days
  a week. This is a private car park and charges apply. The car park is continually
  monitored by parking hosts and CCTV. There are disabled spaces on the level one
  of the car park. If you need help, please speak with a parking host at the car park
  entrance.
- If you have been referred to us from clinic and usually have an ambulance to bring you for your appointment, please contact us as soon as possible on:

Tel: 0151 706 2656

Text phone number: 18001 0151 706 2656

• If you have been referred to us by your family doctor (GP) and require an ambulance contact your family doctor to book the ambulance.

#### **Further appointments**

You may be given a follow up appointment before leaving or one will be sent to you in the post to attend the clinic of the doctor who referred you.

#### Important: If you have:

- Diabetes
- Are taking: (Warfarin tablets, Clopidogrel (plavix) tablets)
- Anticoagulants:
  - Warfarin
  - Heparin/Low molecular weight heparin (including enoxaparin/dalteparin)
  - Dabigatran
  - Rivaroxiban
  - Sinthrome
  - Apixaban
  - Fondaparinux
  - Edoxaban
- Antiplatelet therapy:
  - Clopidogrel (Plavix)
  - Prasugrel (Efient)
  - Ticagrelor (Brilique)
  - Dipyridamole (Persantin) and aspirin

- Are on dialysis.
- Have suffered a heart attack within the last three months.

You must contact the Gastroenterology Unit as soon as you receive this information leaflet.

Tel: 0151 706 2656

Text phone number: 18001 0151 706 2656

#### **Feedback**

Your feedback is important to us and helps us influence care in the future.

Following your discharge from hospital or attendance at your outpatient appointment you will receive a text asking if you would recommend our service to others. Please take the time to text back, you will not be charged for the text and can opt out at any point. Your cooperation is greatly appreciated.

#### **Further information**

If you have any questions or queries, please contact the Gastroenterology Unit during the following hours.

Monday – Thursday	0800 – 2100 hrs
Friday	0800 – 1700 hrs
Saturday/Sunday/BH	0800 – 1600 hrs

Tel: 0151 706 2656/ 2819/2726

Text phone number: 18001 0151 706 2656/2819/2726

Clinic appointment enquiries

Tel: 0151 706 5555

Text phone number: 18001 0151 706 5555

The Emergency Department (A&E) is open 24 hours

Tel: 0151 706 2051/2050.

Text phone number: 18001 0151 706 2051/2050

Hospital: Royal hospital

Tel: 0151 706 2656/ 2819/2726

Text phone number: 18001 0151 706 2656/2819/2726

**Author: Digestive Disease Care Group** 

**Review date: February 2022** 

All Trust approved information is available on request in alternative formats, including other languages, easy read, large print, audio, Braille, moon and electronically.

يمكن توفير جميع المعلومات المتعلقة بالمرضى الموافق عليهم من قبل ائتمان المستشفى عند الطلب بصيغ أخرى، بما في ذلك لغات أخرى وبطرق تسهل قراءتها وبالحروف الطباعية الكبيرة وبالصوت وبطريقة برايل للمكفوفين وبطريقة مون والكترونيا.

所有經信托基金批准的患者資訊均可以其它格式提供,包括其它語言、 易讀閱讀軟件、大字

體、音頻、盲文、穆恩體(Moon)盲文和電子格式,敬請索取。

در صورت تمایل میتوانید کلیه اطلاعات تصویب شده توسط اتحادیه در رابطه با بیماران را به اشکال مختلف در دسترس داشته باشید، از جمله به زبانهای دیگر، به زبان ساده، چاپ درشت، صوت، خط مخصوص کوران، مون و بصورت روی خطی موجود است.

زانیاریی پیّوهندیدار به و نهخوشانه ی له لایه ن تراسته و ه پهسهند کراون، ئهگهر داوا بکریّت له فورماته کانی تردا بریتی له زمانه کانی تر ، ئیزی رید (هاسان خویندنه وه)، چاپی گهوره، شریتی دهنگ، هیّلی موون و نهلیّکتروّنیکی همیه.

所有经信托基金批准的患者信息均可以其它格式提供,包括其它语言、易读阅读软件、大字体、音频、盲文、穆恩体(Moon)盲文和电子格式,敬请索取。

Dhammaan warbixinta bukaanleyda ee Ururka ee la oggol yahay waxaa marka la codsado lagu heli karaa nuskhado kale, sida luqado kale, akhris fudud, far waaweyn, dhegeysi, farta braille ee dadka indhaha la', Moon iyo nidaam eletaroonig ah.