The Royal Liverpool and Broadgreen University Hospitals

NHS Trust



Patient information

Starting Glucophage SR

Directorate of Diabetes and Endocrinology

Royal Liverpool Hospital and Broadgreen Hospital

PIF 1058 V4

Glucophage SR may also be known as Metformin M/R (modified release).

What does SR mean?

SR is an abbreviation for sustained release. This means that the effect of the tablet is slow and even release over a prolonged period of time. In the case of Glucophage SR each tablet will be active for approximately twenty-four hours.

The minimum and maximum dose

The dose will vary from person to person, however the total daily dose ranges from 500 – 2000mg.

How does Glucophage SR work?

Glucophage SR tablets work in two ways. In the first instance it helps to prevent the liver from producing its own glucose. Its other beneficial effect is that it encourages insulin to carry glucose from the blood into the muscle and fat cells more efficiently.

Side effects

As with any medication there can be side effects. For Metformin M/R these can include:

- Upset stomach
- Nausea
- Diarrhoea
- Metallic taste in the mouth.

These effects are reduced if the dose of Metformin is slowly increased once treatment is started and tablets are taken with or immediately after food.

Glucophage (Metformin) is the only medication in its class, this means that unfortunately if a person can not tolerate metformin or metformin M/R there is no other medication in this class that can be used. A different type of diabetes medication may have to be used instead. If you would like more information about different types of diabetes medication speak to your family doctor (GP) or diabetes team.

What effect will Glucophage SR have on my insulin doses or other diabetic tablets?

When Glucophage SR is used it usually has a lowering effect on insulin requirements. This effect is usually seen once two tablets of Glucophage SR 500mg or more are taken on a daily basis.

This may just result in an improvement in your diabetes control, but on some occasions you may need your other diabetic medication or insulin therapy reduced if your blood glucose levels become to low.

Dosage Schedule for Glucophage SR taken once a day.

Glucophage SR should be increased every two weeks.

- Week 1 Take one tablet of Glucophage SR 500mg with your evening meal.
- **Week 3** Take two tablets of Glucophage SR 500mg with your evening meal.
- Week 5 Take three tablets of Glucophage SR 500mg with your evening meal.
- Week 7 Take four tablets of Glucophage SR 500mg with your evening meal.

Alternative dosage Schedule for Glucophage SR taken twice a day

- Week 1 Take one tablet of Glucophage SR 500mg with your evening meal.
- Week 3 Take one tablet of Glucophage SR 500mg with your breakfast and one tablet of Glucophage SR 500mg your evening meal.
- Week 5 Take one tablet of Glucophage SR 500mg with your breakfast and two tablets of Metformin MR 500mg with your evening meal.
- Week 7 Take two tablets of Glucophage SR 500mg with your breakfast and two tablets of Metformin MR 500mg with your breakfast.

Further information

If you want any further information or have any problems with your dosage then please contact the diabetes nurses, your pharmacist or GP (General Practitioner).

The Diabetes Centre Tel: 0151 706 2829

NHS 111 Tel 111 Diabetes UK
The North West Diabetes UK
First floor
The Boultings
Winwick Street
Warrington
WA2 7TT.

Tel: 01925 653 281 Fax: 01925 653 288

E-mail: n.west@diabetes.org.uk Website www.diabetes.org.uk

Author: Diabetes Centre Review date: August 2013

All Trust approved information is available on request in alternative formats, including other languages, easy read, large print, audio, Braille, moon and electronically.

يمكن توفير جميع المعلومات المتعلقة بالمرضى الموافق عليهم من قبل انتمان المستشفى عند الطلب بصبغ أخرى، بما في ذلك لخات أخرى وبطرق تسهل قراءتها وبالحروف الطباعية الكبيرة وبالصوت وبطريقة برايل للمكفوفين وبطريقة مون والكثر ونيا.

所有經信托基金批准的患者資訊均可以其它格式提供,包括其它語言、 易讀閱讀軟件、大字

體、音頻、盲文、穆思體(Moon)盲文和電子格式,敬請索取。

در صورت تمایل میتوانید کلیه اطلاعات تصویب شده توسط اتحادیّه در ایطه با بیماران را به اشکال مختلف در دسترس داشته باشید، از جمله به زبانهای دیگر، به زبان ساده، چاپ درشت، صوت، خط مخصوص گوران، مون و بصورت روی خطی موجود است.

ز انباریی پیوهندیدار به و نهخوشانه ی لهلایهن تراسته و پسمه ند کر اون، نهگهر داوا بکریت له فورماته کانی تر دا بریتی له زمانه کانی تر، نیزی رید (هاسان خویندنه وه)، چایی گهوره، شریتی دهنگ، هیلی موون و نهای کترونیکی هایه.

所有经信托基金批准的患者信息均可以其它格式提供,包括其它语言、易读阅读软件、大字体、音频、盲文、穆恩体(Moon)盲文和电子格式,敬请索取。

Dhammaan warbixinta bukaanleyda ee Ururka ee la oggol yahay waxaa marka la codsado lagu heli karaa nuskhado kale, sida luqado kale, akhris fudud, far waaweyn, dhegeysi, farta braille ee dadka indhaha la', Moon iyo nidaam eletaroonig ah.