

#### Patient information

# **Chalazion / Meibomian Cyst Surgery**

Ophthalmology Department – Aintree Hospital

#### What is a Chalazion?

A common eye lid disorder, a chalazion is an infection or inflammation of a meibomian gland (one of the oil glands) in the upper or lower eyelid, associated with a red, tender swelling and usually develops slowly over several weeks.



#### What Treatment will I need for a Chalazion?

# There are two types of treatment:

#### **Medical**

When the chalazion first appears it is possible to treat with warm compresses and massage.

A small non-infected chalazion usually disappears in one or two months. If it is infected antibiotics may be required.

Occasionally the chalazion may become larger and cause pressure on the eye resulting in some blurring of the vision.

Generally however, leaving the chalazion is harmless. Many chalazion disappear without treatment after a few months.

## **Surgery**

Sometimes medical treatment does not resolve the chalazion. The chalazion may eventually become a hard cyst and require an incision and curettage (drainage).

# What is an incision and curettage?

- This procedure aims to reduce the size of the cyst and is performed as follows:
- A drop of local anaesthetic is instilled into the eye to numb the inside surface of the eyelid.
- A small amount of local anaesthetic is injected into the skin around the cyst.
- A small clamp is placed around the cyst to help prevent bleeding.
- A cut is made into the cyst (on the inside of the eyelid) and the contents of the cyst removed.
- The clamp is removed and pressure is applied to the eye lids to prevent bleeding.
- Antibiotic ointment is put into the eye to prevent infection.
- The eye is padded for a minimum of two hours.

# Are there any risks or adverse reactions involved in having an incision and curettage of chalazion?

• The local anaesthetic injection can sting and be painful, but this enables the surgery to be pain free.

- The surgery may cause bruising of the eyelid.
- Infection at wound site.
- Recurrence / new cysts may develop.
- Further treatment may be required.

## Care after chalazion surgery

You should leave the pad dressing on your eye for at least two hours after surgery.

Do not be alarmed if the eyelids are bruised and swollen. This will settle in a few days. Instil the antibiotic eye ointment three times a day for four days.

- Do not wear contact lenses until your course of eye ointment is finished.
- There will be no postoperative follow up appointment.

#### **Feedback**

Your feedback is important to us and helps us influence care in the future. Following your discharge from hospital or attendance at your outpatient appointment you will receive a text asking if you would recommend our service to others. Please take the time to text back, you will not be charged for the text and can opt out at any point. Your co-operation is greatly appreciated.

# **Further information**

If you have any questions please contact:

Ophthalmic Nurse Specialist: (Monday to Friday 9am-

3pm)

Tel: 0151 529 0196

**Eye Clinic:** 

Tel: 0151 529 0185

**Author: Ophthalmology Department** 

Review date: April 2026

All Trust approved information is available on request in alternative formats, including other languages, easy read, large print, audio, Braille, moon and electronically.

يمكن توفير جميع المعلومات المتعلقة بالمرضى الموافق عليهم من قبل ائتمان المستشفى عند الطلب بصيغ أخرى، بما في ذلك لغات أخرى وبطرق تسهل قراءتها وبالحروف الطباعية الكبيرة وبالصوت وبطريقة برايل للمكفوفين وبطريقة مون والكترونيا.

所有經信托基金批准的患者資訊均可以其它格式提供,包括其它語言、 易讀閱讀軟件、大字

體、音頻、盲文、穆恩體(Moon)盲文和電子格式,敬請索取。

در صورت تمایل میتوانید کلیه اطلاعات تصویب شده توسط اتحادیه در رابطه با بیماران را به اشکال مختلف در دسترس داشته باشید، از جمله به زبانهای دیگر، به زبان ساده، چاپ درشت، صوت، خط مخصوص کوران، مون و بصورت روی خطی موجود است.

زانیاریی پیّوهندیدار به و نهخو شانه ی له لایه ن تراسته و ه پهسهند کراون، نهگه ر داوا بکریّت له فورماته کانی تردا بریتی له زمانه کانی تر، نیزی رید (هاسان خویّندنه وه)، چایی گهوره، شریتی دهنگ، هیّلی موون و نهلیّکترونیکی ههیه.

所有经信托基金批准的患者信息均可以其它格式提供,包括其它语言、易读阅读软件、大字体、音频、盲文、穆恩体(Moon)盲文和电子格式,敬请索取。

Dhammaan warbixinta bukaanleyda ee Ururka ee la oggol yahay waxaa marka la codsado lagu heli karaa nuskhado kale, sida luqado kale, akhris fudud, far waaweyn, dhegeysi, farta braille ee dadka indhaha la', Moon iyo nidaam eletaroonig ah.