

Patient information

Mucous Membrane Pemphigoid (MMP)

Liverpool University Dental Hospital

What are the aims of this leaflet?

This leaflet has been written to help you understand more about mucous membrane pemphigoid (MMP). It tells you what it is, what causes it, what can be done about it, and where you can find more information.

What is MMP?

MMP is the most up-to-date name for this condition. Other names include cicatricial pemphigoid, oral pemphigoid and ocular pemphigoid.

MMP is an uncommon blistering condition which most frequently affects the lining of the mouth and gums.

Other moist surfaces of the body (known as mucous membranes) can also be affected, and these include the surface layers of the eyes, the inside of the nose, the throat and the genitals.

The skin is affected sometimes, usually by a few scattered blisters. MMP usually starts in middle and old age.

Although it is not usually a serious condition in the mouth, the diagnosis of any type of MMP is important as it will alert your specialist to the possibility that the condition may involve your eyes, even if you have no symptoms. Eye involvement (known as ocular cicatricial pemphigoid or ocular MMP) does not occur in all patients but is potentially serious as it may cause scarring and affect your eyesight.

Scarring may also result in significant damage to the throat and the genitals, and may be very serious if the larynx is affected.

What causes MMP?

The cause of MMP is unknown. It is considered to be an autoimmune disease involving antibodies (natural substances important in defending your body against infections) that react with the surface layer of your mouth (or other mucous membranes) causing blisters, which usually break down to leave ulcers.

MMP is not contagious, or due to food allergy.

Is MMP hereditary?

MMP is not an inherited condition.

What are the symptoms of MMP?

- In mild cases, MMP may just involve the mouth causing occasional blisters. These are usually painless until they breakdown to form ulcers that typically cause burning or stinging, especially when eating spicy foods, citrus fruits, drinking hot beverages, alcohol or using mouthwashes that contain alcohol. If your gums are affected, they may become sore, particularly when brushing your teeth.
- In more severe cases the mouth, throat, eyes, nose, genitals or skin may also be affected. If your eyes are affected, they may become red and feel gritty or sore.
- Some people with MMP get a stuffy blocked nose and have blood-tinged mucus or nose bleeds, or notice a change in their voice. Occasionally they may experience some difficulty swallowing.
- If the genitals are affected, blisters or ulcers may appear and tend to persist unless treated.
- There may be scattered skin blisters often on the scalp but occasionally on the face, trunk or limbs.

What does MMP look like?

In the mouth it typically presents as red or ulcerated patches on the inner cheeks, gums or roof of the mouth. Blisters may occur but tend to easily burst and leave ulcers. The gums become red, shiny and ulcerated and may be the only sign of this condition.

Active pemphigoid in the eye will usually appear as a red eye and if not treated may lead to scarring.

Skin blisters may form crusts and eventually heal with a superficial scar.

How is MMP diagnosed?

MMP cannot be diagnosed solely by its appearance as other conditions can look very similar. One or two small samples of tissue are usually taken by the specialist team (under a local anaesthetic), usually inside the mouth (this procedure is known as a biopsy). The diagnosis can then be made by looking at the samples under a microscope and testing them for specific antibodies associated with MMP.

A blood test may also be used to detect these antibodies in the circulation.

You may be referred to an eye specialist (ophthalmologist) who can pick up early signs of pemphigoid involving the eyes, an Ear Nose and Throat (ENT) specialist or a dermatologist depending on your symptoms. This multidisciplinary team care is very important.

Can MMP be cured?

MMP is usually brought under control with systemic medication (tablets or injections). Tablets may be needed for several years but in some patients the condition can go into long term remission and eventually medication may be stopped.

How can MMP be treated?

Your doctor will discuss treatment options and help choose the most suitable medication depending on disease severity, your general health and potential medication side effects.

Topical treatments:

Most patients will be prescribed topical treatments for each affected site:

For the mouth:

- Anaesthetic (analgesic) mouthwashes are available if your mouth becomes sore and are particularly helpful if used before meals. Benzydamine (Diffiam) mouthwash may be helpful.
- Topical steroids which can be applied locally to the mouth are helpful. (most patients however require additional systemic medication (tablets). These are available as mouthwashes, sprays, pastes and small pellets which dissolve in your mouth. Your specialist may also consider the use of topical steroids inside custom made gum shields. Topical steroids can sometimes cause oral thrush which can be easily treated or prevented with topical antifungal treatment.
- If your gums are affected (desquamative gingivitis), it is important that you keep your teeth as clean as possible by regular and effective tooth brushing. If not, a build-up of debris (known as plaque) can make your gum condition worse. Your dentist or dental hygienist will be able to give oral hygiene advice and will arrange for scaling of your teeth as necessary.
- An antiseptic mouthwash or gel may be recommended to help with your plaque control, particularly at times when your gums are sore. Daily hydrogen peroxide mouthwash (Peroxyl) or occasional chlorhexidine (Corsodyl) twice per week are examples. If possible avoid mouthwashes containing alcohol.

For the skin

- Corticosteroid creams or ointments are helpful and will be prescribed by your doctor.

For the eyes

- Treatment will be directed by your ophthalmologist and may include lubricants and topical corticosteroid eye drops although systemic (oral or intravenous) medication is needed for the majority of patients.

Systemic (whole body) treatment

In people who are more severely affected with MMP, or when topical treatment has not helped, systemic treatment (taken by mouth or by injection into a vein) may be required, usually for several months or years. Your specialist will discuss with you risks and benefits of the different drug options available.

Regular blood tests are required, to check for unwanted effects of treatment especially in the first weeks to months of treatment.

- Oral steroids may be used for a few months in high doses, then by some specialists, for longer periods at low doses.
- Other tablets used to treat MMP include dapsone, Sulphamethoxy pyridazine (SMP), sulphapyridine or other antibiotics such as tetracycline combined with nicotinamide.
- Immunosuppressant drugs such as azathioprine, mycophenolate mofetil, methotrexate, cyclophosphamide or rituximab may be needed for severe MMP. These help MMP by suppressing the body's immune system.

What can I do?

- Avoid spicy, acidic or salty foods if these make your mouth sore.
- Have your mouth checked on a regular basis by a dentist, or oral specialist.
- Keep your teeth clean by using a soft brush and small interdental brushes
- Choose a toothpaste with a mild flavour and free from the foaming agent, sodium lauryl sulphate (SLS)
- Stop smoking and reduce your alcohol intake to recommended limits.
- For the nose, nasal douching can be helpful.
- For the eyes, please seek precise advice from your ophthalmologist.

Where can I get more information about MMP?

Web links to detailed leaflets:

- <http://emedicine.medscape.com/article/1062534-overview>
- <http://dermnetnz.org/immune/cicatricial-pemphigoid.html>

Links to patient support groups:

- Pemfriends <https://www.pemfriendsuk.co.uk/>
- The International Pemphigus and Pemphigoid Foundation www.pemphigus.org

Not all the information will be relevant to all patients. For individual advice please see your Oral Medicine specialist.

Feedback

Your feedback is important to us and helps us influence care in the future.

Following your discharge from hospital or attendance at your outpatient appointment you will receive a text asking if you would recommend our service to others. Please take the time to text back, you will not be charged for the text and can opt out at any point. Your co-operation is greatly appreciated.

Further Information

If you need any further information please contact the Liverpool University Dental Hospital

Tel: 0151 706 5060

Text phone number: 18001 0151 706 5060

Author: Reproduced with the Kind Permission of The British & Irish Society of Oral Medicine (BISOM) and the British association of Dermatologists

Review date: March 2027



All Trust approved information is available on request in alternative formats, including other languages, easy read, large print, audio, Braille, moon and electronically.

يمكن توفير جميع المعلومات المتعلقة بالمرضى الموافق عليهم من قبل انتمان المستشفى عند الطلب بصيغ أخرى، بما في ذلك لغات أخرى وبطرق تسهل قراءتها وبالحروف الطباعية الكبيرة وبالصوت وبطريقة برايل للمكفوفين وبطريقة مون والكترونياً.

所有經信托基金批准的患者資訊均可以其它格式提供，包括其它語言、易讀閱讀軟件、大字

體、音頻、盲文、穆恩體 (Moon) 盲文和電子格式，敬請索取。

در صورت تمایل می‌توانید کلیه اطلاعات تصویب شده توسط اتحادیه در رابطه با بیماران را به اشکال مختلف در دسترس داشته باشید، از جمله به زبانهای دیگر، به زبان ساده، چاپ درشت، صوت، خط مخصوص کوران، مون و بصورت روی خطی موجود است.

زانیاری پیوندیدار بهو نمخوشانهی له لایمن تراستهوه پسمند کراون، نهگس داوا بکریت له فورماتهکانی تردا بریتی له زمانهکانی تر، نیزی رید (هاسان خویندنهوه)، چاپی گموره، شریتی دمنگ، هیلی موون و نلیکترونیکي همیه.

所有经信托基金批准的患者信息均可以其它格式提供，包括其它语言、易读阅读软件、大字体、音频、盲文、穆恩体 (Moon) 盲文和电子格式，敬请索取。

Dhammaan warbixinta bukaanleyda ee Ururka ee la oggol yahay waxaa marka la codsado lagu heli karaa nuskhado kale, sida luqado kale, akhris fudud, far waaweyn, dhegeysi, farta braille ee dadka indhaha la', Moon iyo nidaam eletaroonig ah.

