

Patient information

Recurrent Aphthous Stomatitis (Recurrent Mouth Ulcers)

Liverpool University Dental Hospital

What are the aims of this leaflet?

This leaflet has been written to help you understand more about Recurrent Aphthous Stomatitis (RAS). It tells you what it is, what causes it, what can be done about it and where you can find out more about it.

What is recurrent aphthous stomatitis (RAS)?

RAS consists of recurrent bouts of one or more painful, rounded or oval shaped ulcers. Most aphthous ulcers last for ten to fourteen days. It is a common mouth condition affecting up to 20% of the population at any given time. The severity and frequency of RAS tends to decrease with age.

RAS is classified into three types:

Minor ,Major and Herpetiform

- ❖ **Minor** ulcers are the most common type affecting the majority (80%) of people who suffer from mouth ulcers. Minor ulcers occur in crops of about one to five at a given time and are usually 10mm or less in diameter. These usually appear inside the lips and cheeks, on the tongue and sometimes the floor of the mouth. The ulcers tend to last between ten to fourteen days and heal without scarring.
- ❖ **Major** ulcers are less common and affect 10-15% of patients with RAS. Major ulcers tend to be larger and typically greater than 10mm in diameter. They can occur singly or two to three can appear at a time anywhere in the mouth. If the soft palate is involved, swallowing can be difficult. The ulcers may last up to three months and some of the larger ulcers leave a scar when healed.
- ❖ **Herpetiform** ulcers are the least common type affecting 5-10% of cases. The ulcers are small (1-2mm in diameter) and can occur in clusters of more than 20 at a time which can merge to give larger ulcers. They tend to occur in the front of the mouth particularly under the tongue and on the edges of the tongue. They tend to heal within two weeks without scarring. Despite their name, they are not caused by a herpes virus.

What causes recurrent aphthous stomatitis?

The cause of RAS is unknown but there is likely to be an immunological factor involved. There are a number of underlying or precipitating factors which include anaemia, vitamin deficiencies, stress and trauma from sharp teeth, dental braces/fillings or a tooth brush.

Some people who give up smoking develop recurrent mouth ulcers and the reason for this is unclear. Occasionally, RAS can be part of a more widespread disease affecting other parts of the body (e.g. Behçet's disease). Recurrent mouth ulcers are not thought to be infectious.

Is recurrent aphthous stomatitis hereditary?

Nearly half the people who suffer from mouth ulcers have a close relative with the same problem.

What does recurrent aphthous stomatitis look like?

RAS consists of round or oval shaped ulcers with an area of surrounding redness. The base of an ulcer is typically grey or yellow in colour. Size varies according to the type of ulcer and this may range from a few millimetres in diameter to greater than one centimetre.

What are the symptoms of recurrent aphthous stomatitis?

The main complaint is pain. This can be made worse by hot, salty, spicy or hard / abrasive food. Eating and drinking can therefore become difficult. Depending on the site of ulcers, speech can also be affected. The ulcers occur in recurrent bouts, heal and reoccur with varying time intervals. A few people are never free from ulcers.

How is recurrent aphthous stomatitis diagnosed?

The history and clinical appearance of the ulcers are usually sufficient to confirm the diagnosis of RAS. Blood tests are often arranged to check for any underlying cause. A biopsy is occasionally required to rule out other causes of mouth ulceration.

Can recurrent aphthous stomatitis be cured?

There is currently no cure for RAS unless an underlying cause is found and corrected. Treatment aims to relieve the painful symptoms associated with RAS. The frequency and severity of RAS tends to decrease with age.

How can recurrent aphthous stomatitis be treated?

Treatment for RAS aims to relieve discomfort, prevent or reduce secondary infection and encourage healing.

- Topical corticosteroids are the main treatment for RAS. They can be applied locally to the mouth and are effective for most patients. These are available as mouthwashes, sprays, and small dissolvable pellets.
- Anaesthetic (analgesic) mouthwashes, sprays or 'over-the-counter' sugar free throat lozenges can be used if your mouth becomes sore and are particularly helpful if used before meals.
- Use of an antiseptic alcohol-free mouthwash, spray or gel (e.g. chlorhexidine gluconate) may be recommended to help reduce any secondary infection and control plaque levels on teeth if tooth brushing is difficult or uncomfortable.
- Tetracycline mouthwashes may be of value for some types of RAS.

- Covering agents work by forming a mechanical barrier against secondary infection and further mechanical irritation. These are available as pastes and soluble pellets for application to ulcers.
- Severe cases of RAS may require treatment with a short course of systemic corticosteroids (i.e. taken in tablet form). Long-term treatment with these drugs is not recommended because of the potential side effects.
- Other types of oral (systemic) therapy are reserved for severe cases of RAS (especially the major type) and various drugs have been used as management options to suppress the ulcers by altering the body's immune system. These can be associated with a number of side effects which should be discussed with your specialist. Regular blood tests are required when taking most of these drugs, particularly during the early stages of treatment.

What can I do?

- Spicy, acidic, salty or hard/abrasive foods (e.g. toast and crisps) should be avoided if these make your mouth sore.
- To avoid nutritional deficiencies, ensure that you eat a varied diet.
- It is important to have a high standard of oral hygiene.
- Visit your dentist for routine care and for help with sharp teeth or broken fillings that may trigger your mouth ulcers.
- Additives in toothpastes, such as sodium lauryl sulphate, may aggravate your ulcers.
- Any single ulcer that persists for longer than three weeks despite treatment should be examined by your dentist (or doctor) who may wish to refer you for a specialist opinion and possible biopsy.

Where can I get more information about recurrent aphthous stomatitis?

<http://www.dermnetnz.org/site-age-specific/aphthae.html>

<http://www.mayoclinic.com/health/canker-sore/DS00354>

Feedback

Your feedback is important to us and helps us influence care in the future.

Following your discharge from hospital or attendance at your outpatient appointment you will receive a text asking if you would recommend our service to others. Please take the time to text back, you will not be charged for the text and can opt out at any point. Your co-operation is greatly appreciated.

Further Information

**If you need any further information please contact the Liverpool University Dental Hospital
Tel: 0151 706 5060**

Author: This information has been reproduced with the kind permission of the British and Irish Society of Oral Medicine (BISOM)
Review date: March 2027

All Trust approved information is available on request in alternative formats, including other languages, easy read, large print, audio, Braille, moon and electronically.

يمكن توفير جميع المعلومات المتعلقة بالمرضى الموافق عليهم من قبل انتمان المستشفى عند الطلب بصيغ أخرى، بما في ذلك لغات أخرى وبطرق تسهل قراءتها وبالحروف الطباعية الكبيرة وبالصوت وبطريقة برايل للمكفوفين وبطريقة مون والإلكترونية.

所有經信托基金批准的患者資訊均可以其它格式提供，包括其它語言、易讀閱讀軟件、大字體、音頻、盲文、穆恩體 (Moon) 盲文和電子格式，敬請索取。

در صورت تمایل می‌توانید کلیه اطلاعات تصویب شده توسط اتحادیه در رابطه با بیماران را به اشکال مختلف در دسترس داشته باشید، از جمله به زبانهای دیگر، به زبان ساده، چاپ درشت، صوت، خط مخصوص کوران، مون و بصورت روی خطی موجود است.

زانیاری پنهانیدار بهو نه‌خوشانه‌ی له‌لایمن تراسته‌وه پسه‌ند کراون، نه‌گس داوا بکرنیت له فورمات‌ه‌کانی تردا بریتی له زمانه‌کانی تر، نیزی رید (هاسان خویندنه‌وه)، چاپی گه‌وره، شریتی ده‌نگ، هیلای موون و نه‌لنیکترونیکی هیه.

所有经信托基金批准的患者信息均可以其它格式提供，包括其它语言、易读阅读软件、大字体、音频、盲文、穆恩体 (Moon) 盲文和电子格式，敬请索取。

Dhammaan warbixinta bukaanleyda ee Ururka ee la oggol yahay waxaa marka la codsado lagu heli karaa nuskhado kale, sida luqado kale, akhris fudud, far waaweyn, dhegeysi, farta braille ee dadka indhaha la', Moon iyo nidaam eletaroonig ah.