

Patient information

Surgical Removal of Teeth or Roots

Special Care Dentistry Department – Liverpool University Dental Hospital

Your dentist has advised you that you will need your tooth (or roots) surgically removed. This is known as a “minor oral surgical procedure” or “M.O.S”. This leaflet has been designed to improve your understanding of this treatment and contains answers to many of the commonly asked questions. If you have any other questions, or would like further explanation, please ask a member of the clinical staff.

The problem

The tooth or roots cannot be taken out simply with forceps, there can be many reasons for this:-

- If teeth or roots are not completely visible in the mouth they cannot be removed using ordinary dental forceps.
- The roots maybe angled in different directions so the tooth may need to be divided.
- The gum may have “grown over” the root.
- The crown of the tooth may be completely decayed, so it “breaks off” the root during extraction.

Why do I need treatment?

Retained roots and broken down teeth can cause pain and may also become infected causing abscesses.

What does surgical removal involve?

- This involves making a small cut in the gum, so this can be lifted off the bone – this is called a “flap”.
- It may be necessary to remove a small amount of bone from around the tooth, (roots), which may need to be divided before removal.
- The gum is then put back into place with stitches.
- In the majority of cases the stitches are dissolvable and take around two weeks to disappear.

Are there any alternative treatments available?

The only alternative is to leave the teeth (roots) as they are and the problems will continue.

What will happen if I don't have any treatment?

- The tooth / roots will continue to cause problems, which may become more severe.
- Infection is likely to recur and result in swelling and pain.
- Serious complications, such as infection of the jaw bone (osteomyelitis) or abscess formation in the neck rarely occur but can be life threatening.

What type of anaesthetic is used?

A number of options are available depending on how difficult the root or tooth is to remove. Commonly, a local anaesthetic is used.

- **Local anaesthetic** - this is an injection into the gum surrounding the tooth or roots, rather similar to that you may have had for a dental filling. The injection takes a couple of minutes to numb the area and means that you will feel no pain while the surgery is carried out.
- Like all medicines, local anaesthetics may sometimes cause side effects, as well as the effects that are needed. You may experience dizziness, blurred vision, drowsiness and occasionally loss of consciousness.
- Serious side effects are rare and include fits, low blood pressure, slowed breathing and changes in heartbeat, which may be life-threatening. If you have concerns about any of these effects you should discuss them with your dentist.
- **Local anaesthetic and intravenous sedation** – in addition to a local anaesthetic injection you can be given an injection into your arm. This makes you feel relaxed and less aware of the procedure. Further information is available in PIF leaflet number 641.
- **General anaesthetic** – rarely it may be necessary to carry out surgical extraction of teeth or roots under a “day case” general anaesthetic, i.e. although you are put to sleep completely you will be able to go home on the same day as surgery. Further information is available in the Royal College of Anaesthetists patient information ‘You and your Anaesthetic’.

What can I expect after the operation?

When the local anaesthetic wears off a few hours after surgery you will have some discomfort. Your surgeon will give you advice about painkillers. Occasionally, it is necessary to take a course of antibiotics. The discomfort is usually worse for the first few days although it may take a couple of weeks to completely disappear. You may need a few days off work, during which time you should avoid strenuous exercise.

Some swelling can occur both inside and outside the mouth after surgery. This is usually most noticeable for about two days.

There may also be bruising due to bruising under the skin in this area.

It is important to keep the site of surgery as clean as possible for the first few weeks after surgery. You will be given full instructions about the care of your mouth after the surgery is completed.

What are the possible problems?

It is unusual for the area to bleed excessively after surgery, but if this happens applying pressure over the area for at least 10 minutes with a rolled up handkerchief or swab can usually stop it. If the bleeding does not stop you will need to contact the department (or on-call dentist/ Emergency Department (A+E) outside of working hours).

In some parts of the mouth there are nerves which are close to the roots of the teeth, there is always a slight risk of “bruising” to these nerves if surgery is carried out in these areas – this can result in some temporary “numbness” afterwards –your surgeon will tell you if you are particularly at risk of this and show you on your X-ray where the nerves are situated.

Some upper teeth, particularly molars, can be close to the maxillary sinus and a hole into the sinus can occur, following their removal – again your dentist will show you this on the x-ray if you are at risk. If this complication does occur you may need to have a small surgical repair carried out and avoid blowing your nose for two weeks afterwards.

Access to Emergency advice

The Liverpool University Dental Hospital is open 9am to 5pm on weekdays (excluding public holidays). If you have problems outside of these times e.g. severe pain, not relieved by pain killers, mild post-operative bleeding or facial swelling please contact the dental urgent care line on

Tel: 0161 476 9651

Text phone number: 18001 0161 476 9651

If you have problems with uncontrolled bleeding, severe facial swelling or feel systemically unwell following dental treatment please contact your local Emergency Department (A+E) for advice.

Feedback

Your feedback is important to us and helps us influence care in the future.

Following your discharge from hospital or attendance at your outpatient appointment you will receive a text asking if you would recommend our service to others. Please take the time to text back, you will not be charged for the text and can opt out at any point. Your co-operation is greatly appreciated.

Further Information

Special Care Dentistry Department

Tel: 0151 706 5530/5235

Text phone number: 18001 0151 706 5530/5235

All Trust approved information is available on request in alternative formats, including other languages, easy read, large print, audio, Braille, moon and electronically.

يمكن توفير جميع المعلومات المتعلقة بالمرضى الموافق عليهم من قبل انتمان المستشفى عند الطلب بصيغ أخرى، بما في ذلك لغات أخرى وبطرق تسهل قراءتها وبالحروف الطباعية الكبيرة وبالصوت وبطريقة برايل للمكفوفين وبطريقة مون والإلكترونية.

所有經信托基金批准的患者資訊均可以其它格式提供，包括其它語言、易讀閱讀軟件、大字

體、音頻、盲文、穆恩體（Moon）盲文和電子格式，敬請索取。

در صورت تمایل می‌توانید کلیه اطلاعات تصویب شده توسط اتحادیه در رابطه با بیماران را به اشکال مختلف در دسترس داشته باشید، از جمله به زبانهای دیگر، به زبان ساده، چاپ درشت، صوت، خط مخصوص کوران، مون و بصورت روی خطی موجود است.

زانیاری پنیوندیدار بهو نهخوشانهی له‌لایمن تراستهوه پسمند کراون، نهگس داوا بکریت له فورمات‌هکانی تردا بریتی له زمانه‌کانی تر، نیزی رید (هاسان خویندنهوه)، چاپی گه‌وره، شریتی دهنگ، هیلای موون و نهلیکترونیکی هیه.

所有经信托基金批准的患者信息均可以其它格式提供，包括其它语言、易读阅读软件、大字体、音频、盲文、穆恩体（Moon）盲文和电子格式，敬请索取。

Dhammaan warbixinta bukaanleyda ee Ururka ee la oggol yahay waxaa marka la codsado lagu heli karaa nuskhado kale, sida luqado kale, akhris fudud, far waaweyn, dhegeysi, farta braille ee dadka indhaha la', Moon iyo nidaam eletaroonig ah.